


Sub
Bⁱ

~~(±)-Benzhydrylsulfinylacetamide~~

5 SUP The present invention relates to the levo-
rotatory derivative of benzhydrylsulfinylacetamide,
the method for its preparation and its use in therapy,
especially as an antidepressant and stimulant for the
central nervous system (CNS), the said derivative being
useful in particular in the treatment of hypersomnia
on account of its arousing effects.

10 It is known that French Patent no. 78 05 510
(publication no. FR-B-2 385 693) describes the racemate
(±)-benzhydrylsulfinylacetamide, which has the code
number CRL 40 476 and the following structural formula:


T0020X



15 PS as a product (see Example 1 of the said French patent)
and as a stimulant for the central nervous system (CNS).

17 P It is also known that, in Patent Document EP-A
0 097 071, the neuropsychopharmacological properties of
the racemate were compared with those of the analogs of
the formula:

20 T0021X



Pl in which:

EPI X_1 and X_2 , which can be identical or different, each represent H, Cl or F,

5 ~~EPI~~ Z_1 represents CH_3 , CH_2CH_3 , $CH(CH_3)_2$ or $C(CH_3)_3$, it also being possible for Z_1 to represent a hydrogen atom when at least one of the symbols X_1 and X_2 is different from H, and

~~EPI~~ Z_2 represents H, it being possible for NZ_1Z_2 , considered together, to represent a piperidino or
10 morpholino group,

~~R~~ which act on the CNS as sedatives in some cases and as stimulants in others (see especially Table I on page 3 and Table IV on page 4 of the said European patent document).

15 ρ It has now been found that the levorotatory compound $(-)$ -benzhydrylsulfinylacetamide (Code no.: CRL 40 982) has³¹ valuable therapeutic properties compared with the racemate (\pm) -benzhydrylsulfinylacetamide (Code no.: CRL 40 476)³⁵ and with the dextrorotatory compound
20 $(+)$ -benzhydrylsulfinylacetamide (Code no.: CRL 40 983). Surprisingly, it has been found that the metabolism of the levorotatory compound in the organism is different from that of the racemate and the dextrorotatory compound and that the levorotatory compound is particularly
25 valuable in the treatment of hypersomnia and Alzheimer's disease.

According to the invention, a novel industrial product is recommended which is useful in therapy and belongs to the family of the benzhydrylsulfinyl derivatives, the said product being $(-)$ -benzhydrylsulfinyl-
30 acetamide.³¹

This levorotatory compound cannot be prepared by isolation from the corresponding racemic amide. However, it can be prepared by chemical synthesis from a
35 precursor of the amide, according to a method known per

se, by the application of conventional reaction mechanisms.

The method of preparation recommended according to the invention consists in:

5 PO(1°) reacting (+)-benzhydrysulfinylacetic acid with (-)- α -methylbenzylamine^{31 60} to give the (-)-benzhydrysulfinylacetate of (-)- α -methylbenzylamine^{31 60} (the reaction advantageously being carried out in the presence of a small excess of amine relative to the stoichiometric conditions, and more particularly with a molar ratio
10 amine/acid of between 1.02/1 and 1.15/1 and preferably of between 1.05/1 and 1.10/1),

PO(2°) converting the resulting (-)-benzhydrysulfinylacetate salt of (-)- α -methylbenzylamine^{31 60} to (-)-benzhydrysulfinylacetic acid^{31 60} [the conversion advantageously being carried out by hydrolysis in an acid medium, the solvent being warm water (especially water at 30-45°C)], and

PO(3°) subjecting the resulting (-)-benzhydrysulfinylacetic acid^{31 60} to an amidation reaction with gaseous ammonia.

P The amidation of stage (3°) is advantageously carried out in 2 steps, namely:

PO(3a) esterification of the (-)-benzhydrysulfinylacetic acid to a lower alkyl (-)-benzhydrysulfinylacetate [the lower alkyl being C₁₋₃, especially isopropyl, ethyl^{31 60} or methyl (preferably ethyl and particularly preferably methyl)], followed by

PO(3b) transamidation of the resulting lower alkyl (-)-benzhydrysulfinylacetate with NH₃ (the transamidation reaction preferably being carried out in a lower alcohol and particularly preferably in the alcohol corresponding to the alkyl group of the ester obtained in stage 3a), a stream of NH₃ being passed into the
35 reaction medium).

4

(+)-Benzhydrylsulfinylacetic acid is a known substance³⁵ which is described as a synthesis intermediate in Patent Document FR-B-2 326 181 (m.p. (inst.) = 164³¹ - 165°C).

5 According to the invention, a therapeutic composition is recommended which contains (-)-benzhydrylsulfinylacetamide as the active ingredient³¹, in association with a physiologically acceptable excipient. Of course, in a composition of this type, the said (-)³¹ benzhydrylsulfinylacetamide is present in a pharmaceutically effective amount.

It is also recommended to use this levorotatory compound to obtain, on the one hand, an arousing drug to be used in human therapy for hypersomnia, and, on 15 the other hand, a stimulating drug, and in particular a drug for inhibiting aphasia and ideomotor apraxia, to be used in therapy for Alzheimer's disease.

DEP Further advantages and characteristics of the invention will be understood more clearly from the 20 following description of (i) preparation examples and (ii) results of comparative neuropsychopharmacological tests. These data, which in no way imply a limitation, are given by way of illustration.

CL PREPARATION I

25 Preparation of (-)-benzhydrylsulfinylacetamide

CL (Example 1; Code no.: CRL 40 982)

P(a) (-)-Benzhydrylsulfinylacetate of (-)- α -methyl-³¹benzylamine^{31 60}

30 13 g (0.108 mol) of (-)- α -methylbenzylamine are added to a suspension of 27.4 g^{31 60} (0.1 mol) of (+)³⁵ benzhydrylsulfinylacetic acid (m.p. (inst.) = 164³¹ - 165°C; Code no.: CRL 40 467) in 500 ml of water; the mixture is filtered hot, the filtrate is cooled and the product is filtered off and recrystallized twice from 300 ml of

S

water to give 17 g (yield: 42%) of the (-)-benzhydrylsulfinylacetate of (-)- α -methylbenzylamine.

M.p. (inst.) = $148\frac{31}{11}$ $150\frac{60}{60}$ °C.

P(b) (-)-Benzhydrylsulfinylacetic acid

5 31 The (-)-benzhydrylsulfinylacetate of (-)- α -methylbenzylamine (17 g) obtained in this way is dissolved in 800 ml of warm water ($30\frac{14}{14}$ - $40\frac{60}{60}$ °C) and then acidified with 7 ml of concentrated hydrochloric acid (12 N HCl, d = 1.19 g/cm³). The mixture is filtered
10 cold and the precipitate is washed with water and dried to give the expected (-)-benzhydrylsulfinylacetic acid with a yield of about 31 100%.

P M.p. (inst.) = $185\frac{14}{14}$ - $188\frac{14}{14}$ °C.

14 $20\frac{60}{60}$ °C = 31 -35° (in a 1% solution in CH₃OH).

15 P(c) Methyl (-)-benzhydrylsulfinylacetate

P A suspension of 16.45 g (0.06 mol) of (-)-benzhydrylsulfinylacetic acid in 300 ml of water is treated at 20°C with 16.8 g (0.2 mol) of sodium bicarbonate and 18.8 ml (0.21 mol) of methyl sulfate, with
20 stirring, the mixture is stirred for 16 to 18 hours at 20°C and filtered and the material on the filter is washed with water and dried to give methyl (-)-benzhydrylsulfinylacetate with a yield of 85%.

P M.p. (inst.) = $109\frac{14}{14}$ - $110\frac{14}{14}$ °C.

25 14 $20\frac{60}{60}$ °C = 31 -22.5° (in a 4% solution in CH₃OH).

P(d) CRL 40 982

P A dry stream of NH₃ gas is passed at room temperature into a solution of 100 ml of methanol containing 8.6 g (0.03 mol) of methyl (-)-benzhydrylsulfinylacetate. NH₃ introduced in this way is
30 reacted with the solution for 5 h, with stirring. The methanol is evaporated off, the evaporation residue is taken up in ether and the product is filtered off and recrystallized from ethanol to give CRL 40 982 with an
35 overall yield of 32%. This product is in the form of

white crystals which are soluble in alcohols and acetone and insoluble in water and ether.

M.p. (inst.) = 153-154°C.

$\alpha_D^{20} = -20^\circ$ (in a 2% solution in CH₃OH).

5 CL PREPARATION II

Preparation of (+)-benzhydrylsulfinylacetamide

(Comparison product CP1; Code no.: CRL 40 983)

The following are obtained successively using the procedure indicated in Preparation I above but replacing the (-)- α -methylbenzylamine with (+)- α -methylbenzylamine.

(a) the (+)-benzhydrylsulfinylacetate of (+)- α -methylbenzylamine;

M.p. (inst.) = 148-150°C;

15 (b) (+)-benzhydrylsulfinylacetic acid;

M.p. (inst.) = 190-191°C,

$\alpha_D^{20} = +45^\circ$ (in a 1% solution in CH₃OH);

(c) methyl (+)-benzhydrylsulfinylacetate;

M.p. (inst.) = 109-110°C,

20 $\alpha_D^{20} = +22.2^\circ$ (in a 4% solution in CH₃OH);

and then

(d) CRL 40 983;

M.p. (inst.) = 153-154°C,

$\alpha_D^{20} = +22^\circ$ (in a 2% solution in CH₃OH).

25 The comparative tests which were undertaken with the levorotatory derivative according to the invention (Ex. 1; Code no.: CRL 40 982), the dextro-rotatory derivative (CP1; Code no.: CRL 40 983) and the corresponding racemate (CP2; Code no.: CRL 40 476) have been summarized below. Unless indicated otherwise, the 3 products studied in these tests were administered intraperitoneally as a suspension in an aqueous solution of gum arabic, in a volume of 20 ml/kg to male mice and in a volume of 5 ml/kg to male rats.

7

ρ_A $\bar{13}$ TOXICITY

ρ In male mice, the LD_0 (maximum non-lethal dose) by intraperitoneal administration is found to be greater than or equal to 512 mg/kg for the dextrorotatory compound and the racemate, whereas the LD_{30} of the levo-rotatory compound is of the order of about 512 mg/kg.

ρ In summary, CRL 40 982 is more toxic than CRL 40 483 (CP1) and CRL 40 476 (CP2). The fact that the toxicity of CRL 40 982 is greater than that of the other two products does not present a problem since the levo-rotatory compound still has a sufficiently wide useful range of non-lethal concentrations. Here, the fact that CRL 40 982 is more toxic than the other two products indicates that it is more active.

ρ_{B13} BEHAVIOR IN RATS

ρ In male mice, CRL 40 982, CRL 40 983 and CRL 40 476 have stimulant effects; in male rats, on the other hand, it is found that CRL 40 982 and CRL 40 983 do not have stimulant effects while the racemate (CRL 40 476) (i) is a stimulant and (ii) has a mydriatic action at all the doses used, the levorotatory and dextrorotatory isomers being devoid of this mydriatic action when administered on their own:

ρ at a dose of 128 mg/kg, CRL 40 476 causes excitation with an increase in the fear reaction for 2 h, exophthalmos for 1 h and mydriasis for 1 to 2 h;

ρ at a dose of 32 mg/kg, CRL 40 476 causes excitation (transient, lasting 0.5 h) with an increase in the fear reaction for 1 h, exophthalmos for 0.5 - 1 h and mydriasis for 1, 2 h;

ρ at a dose of 8 mg/kg, CRL 40 476 causes exophthalmos for 0.5 - 1 h and mydriasis for 0.5 h;

ρ at a dose of 2 mg/kg, CRL 40 476 induces transient mydriasis appearing 1 h after administration,

8

whereas
P₁₃ at doses of 64 mg/kg, 16 mg/kg, 4 mg/kg and 1 mg/kg,
CRL 40 982 and CRL 40 983 cause behavior, reactivities
and variations in the rectal temperature and pupil
5 diameter which are substantially comparable to those
of the control group.

P₁₃ C MOTOR ACTIVITY IN MICE


P The mice (6 per dose, 18 control animals)
receive CRL 40 476, CRL 40 982 or CRL 40 983 four hours
10 before being placed in an actimeter, where their motility
is recorded for 30 minutes. It is found that, at doses
of 128 mg/kg and to a lesser extent 64 mg/kg, the three
substances used cause an increase in the motor activity
four hours after their administration. However, the
15 hyperactivity induced by CRL 40 982 and CRL 40 983
reaches a level and a degree of statistical significance
which are greater than those due to CRL 40 476,
especially at the highest dose used (128 mg/kg).

In summary, under the experimental conditions
20 (intraperitoneal administration of the substances four
hours before the test), CRL 40 982 shows a stimulant
effect equal in intensity to that observed with CRL 40
983, while the hyperactivity induced by CRL 40 476 is
less than that obtained with either CRL 40 982 or CRL
25 40 983.

P₁₃ D PHARMACOKINETIC STUDY

P In the organism, CRL 40 476 is partially con-
verted to (±)-benzhydrylsulfinylacetic acid (CRL 40 467)
of the structural formula: PS

30



PS which is used as the starting material for the synthesis

9

of the optical isomers CRL 40 982 and CRL 40 983.

Now, this metabolite is found to be inactive. In fact, CRL 40 467, administered intraperitoneally to male mice at doses of 1024 mg/kg, 512 mg/kg, 256 mg/kg, 128 mg/kg and 64 mg/kg, induces the appearance of brief sedation (with a duration less than or equal to about 50-60 minutes) and does not cause the death of any of the animals treated. Its neuropsychopharmacological study did not reveal any psychotropic activity.

To distinguish CRL 40 982 from the racemic and dextrorotatory compounds, a kinetic study of the metabolism was undertaken on dogs (group of four animals). In a randomized crossover procedure, each animal received the following at the rate of one oral administration per week: 2 administrations of CRL 40 476 (for assessment of the variation within individuals), 1 administration of CRL 40 982 and 1 administration of CRL 40 983. Following each of these administrations, the kinetics of the CRL 40 476 and CRL 40 467 present in the plasma were determined (without looking to see whether these two products were in the racemic, levorotatory and/or dextrorotatory form, because of the difficulties associated with determining the optical rotation of each of the optical isomers in a biological medium).

The dose administered for each of the test substances CRL 40 476, CRL 40 982 and CRL 40 983 was 30 mg/kg.

After administration of CRL 40 476, the said CRL 40 476 and its metabolite, CRL 40 467, are found in the plasma.

After administration of CRL 40 982, the following are found in the plasma: the said CRL 40 982, which will be characterized and determined as being CRL 40 476 by way of convenience, in view of what has been

10

said above, and a metabolite which will be characterized and determined as being CRL 40 467.

Likewise, after administration of CRL 40 983, the 2 corresponding products in the plasma will be characterized and determined as being CRL 40 476 and CRL 40 467.

The curves of the plasma concentrations of the said CRL 40 476 and CRL 40 467 as a function of time are plotted from time $T = 0$ to time $T = +9$ h after administration of the CRL 40 476, CRL 40 982 and CRL 40 983. The areas under the curves (AUC_{0-9h}) are then calculated. The results obtained are recorded in Table I below. They show that:

(a) after administration of CRL 40 476 and CRL 40 983, the AUC_{0-9h} of CRL 40 476 are not statistically different, whereas, after administration of CRL 40 982, the AUC_{0-9h} of CRL 40 476 is approximately twice the AUC_{0-9h} of CRL 40 476 which each result from the administration of CRL 40 476 and CRL 40 983; and

(b) the quantity of CRL 40 467 produced after administration of CRL 40 983 is very large ($83.12 \text{ mg.l}^{-1} \cdot \text{h}$), whereas the quantity produced after administration of CRL 40 982 is very small ($8.69 \text{ mg.l}^{-1} \cdot \text{h}$).

The value of CRL 40 982 according to the invention is in the fact that only a small proportion of this product is converted to inactive CRL 40 467, whereas a very high proportion of the corresponding dextrorotatory derivative is metabolized to CRL 40 467. In summary, the levorotatory compound CRL 40 982 has a better bio-availability than the racemic compound CRL 40 476 and the dextrorotatory compound CRL 40 983, in view of the small quantity of inactive metabolite which it produces in the organism.

These pharmacokinetic results were confirmed on rabbits and mice. An immunostimulant effect was also

observed in vitro for CRL 40 982.

P E - CLINICAL TRIALS

5 ¹³ P In human clinical trials, it was found that the elimination half-life of CRL 40 982 is relatively long (about 10 h), making it possible to obtain good results on adults with 1 to 2 administrations per day.

10 In the course of the clinical trials, CRL 40 982 was found to act in the short term as a purely hypnogenic, antiarousing substance and in the long term as an arousing substance useful for hypersomnia. Furthermore, in both the short and long term, CRL 40 982 was shown to be particularly active towards the symptoms of dementia and loss of memory (especially in the elderly).

15 Administered once or twice a day in the form of tablets or gelatin capsules each containing 50 to 100 mg of CRL 40 982, this product has a different stimulant neuropsychopharmacological profile from that of the amphetamines and tricyclic antidepressants and is useful for depressions, hypersomnia and in particular Alzheimer's disease (improvement of the symptoms of dementia, memory disorders, aphasia and ideomotor apraxia).

12

TABLE I
PHARMACOKINETIC STUDY ON DOGS
MEASUREMENT OF THE AREAS UNDER THE CURVES (AUC_0^{+9h})

Product administered	Code no.	AUC_0^{+9h} CRL 40 476 (a)	AUC_0^{+9h} CRL 40 467 (a)
CP2	CRL 40 476	46.76 ± 6.95	35.12 ± 6.93
Ex. 1	CRL 40 982	97.22 ± 12.58	8.69 ± 1.22
CP1	CRL 40 983	50.94 ± 8.77	83.12 ± 21.66

Notes

(a) in $mg.l^{-1}.h$

* statistically significant difference ($p < 0.1$)

** statistically very significant difference ($p < 0.01$)

n.s. statistically non-significant difference